

Spontaneous Orbifold Symmetry Breaking and Generation of Mass Hierarchy

Israel Quiros*

Departamento de Fisica. Universidad Central de Las Villas. Santa Clara. CP: 54830 Villa Clara.

Cuba

(February 1, 2008)

Abstract

A very simple mechanism is proposed that stabilizes the orbifold geometry within the context of the Randall-Sundrum proposal for solving the hierarchy problem. The electro-weak TeV scale is generated from the Planck scale by spontaneous breaking of the orbifold symmetry.

04.50.+h, 98.80.Cq

Typeset using REVTeX

*israel@uclv.etecsa.cu

M-theory represents the most remarkable theoretical success of the end of the millennium. The moduli space of this theory contains all five, anomaly free, ten-dimensional superstring theories and the eleven-dimensional supergravity. In this context the most successful cosmological construction is that of Horava and Witten [1]. Compactification of the Horava-Witten theory on an eleven-dimensional orbifold $R^{10} \times S^1/Z_2$ to four dimensions on a deformed Calabi-Yau manifold yields that the resulting theory has $N = 1$ supersymmetry. This implies, in turn, that the early universe has undergone a phase where it was five-dimensional [2]. In Ref. [3] an effective five-dimensional theory was derived by direct compactification of the Horava-Witten theory on a Calabi-Yau space. A static solution to the field equations of this theory exists that may be interpreted as a pair of parallel three-branes that are located at the fixed points of the circle that represent the boundaries of the orbifold S^1/Z_2 .

A set-up with two three-branes located at the boundaries of a five-dimensional AdS_5 spacetime has been used recently by Randall and Sundrum to address the hierarchy problem [4]. They proposed an scenario where the metric is not factorizable. The four-dimensional metric, in this scenario, is multiplied by a "warp" factor which is a function of the additional dimension

$$ds^2 = e^{-2k|y|} \eta_{mn} dx^n dx^m + dy^2, \quad (1)$$

where k is a scale of order the Planck scale, x^a are the usual 4-d coordinates and $0 \leq y \leq \pi\lambda$ is the coordinate of the extra dimension. This line-element is consistent with orbifold symmetry ($y \rightarrow -y$) and with four-dimensional Poincare invariance [4]. Randall and Sundrum have shown that this metric is a solution to Einstein's equations in a simple set-up with two three-branes and appropriate cosmological terms. The two three-branes are located at the orbifold fixed points $y = 0$ and $y = \pi\lambda$. These represent the boundaries of the five-dimensional spacetime.

Working out the consequences of the localized energy density inherent to the brane set-

up, Randall and Sundrum found a new solution to the hierarchy problem. In the Randall-Sundrum proposal four-dimensional mass scales are related to five-dimensional input mass parameters and the "warp" factor, $e^{-2k|y|}$. This small exponential factor is the source of the large hierarchy between the observed Planck and weak scales [4]. In this set-up the relation between the Planck scale and the fundamental scale is found to be

$$M_{Pl}^2 = 2M^3 \int_0^{\pi\lambda} dy e^{-2k|y|} = \frac{M^3}{k}(1 - e^{-2k\pi\lambda}), \quad (2)$$

and M_{Pl}^2 is a well-defined value even in the limit $\lambda \rightarrow \infty$, in contrast to the product-space expectation that $M_{Pl}^2 = M^3\lambda\pi$. By taking the second "regulator" brane at infinity and considering the coordinate $y = 0$ to be the location of the Planck brane, one can derive [5]

$$M_{Pl}^2 = 2M^3 \int_0^\infty dy e^{-2k|y|} = \frac{M^3}{k}, \quad (3)$$

so that if M and k are of order $M_{Pl} = 10^{19} GeV$, the graviton zero mode is coupled correctly to generate four-dimensional gravity. For a brane located a distance y_0 from the Planck brane we have $M_{Pl} e^{-k|y_0|} \sim TeV$, i. e., the electroweak scale (of order TeV) is reproduced by physics confined to the brane located a distance y_0 from the brane where the graviton is localized. The generation of this hierarchy requires an exponential of order 30. The advantage of taking a five-dimensional spacetime with infinite extension in the y -direction is that one has a better chance of addressing issues such as the cosmological constant problem and black-hole physics [5].

In this letter we further exploit the ideas of Randall and Sundrum. We present a very simple mechanism of orbifold geometry stabilization that allows determining the location of the TeV brane. It is literally a mechanism of spontaneous orbifold symmetry breaking. Our starting point is a generalization of the Randall-Sundrum 5-d line-element in the form

$$ds^2 = e^{-2k\sigma(y)} g_{mn}(x) dx^m dx^n + e^{l\gamma(y)+h(x)} dy^2, \quad (4)$$

where l is an arbitrary constant factor, σ and γ are arbitrary functions of the additional

y -coordinate, while the four-dimensional metric g_{ab} and h are functions of the familiar 4-d coordinates x^a . For our purposes it suffices to study a general five-dimensional spacetime without fixing the matter content. Neither standard model nor hidden matter sectors are specified. Therefore, we do not pretend to give any realistic particle picture, but a general five-dimensional gravity set-up. The effective 5-d action we shall study is,

$$S = \int d^5x \sqrt{-G} (2M^3 R - \Lambda), \quad (5)$$

where G is the determinant of the five-dimensional metric G_{AB} ($A, B = \overline{0, 4}$), R is the Ricci scalar obtained from the 5-d Ricci tensor R_{AB} ($R = G^{MN} R_{MN}$) and Λ is a cosmological constant that plays the role of a vacuum energy density in the five-dimensional spacetime. The field equations derivable from (5) are

$$R_{AB} - \frac{1}{2} G_{AB} R = -\frac{\Lambda}{4M^3} G_{AB}. \quad (6)$$

These equations can be split into the following set of equations:

$$R_{ab} - \frac{1}{2} g_{ab} e^{-2k\sigma} R = -\frac{\Lambda}{4M^3} e^{-2k\sigma} g_{ab}, \quad (7)$$

$$R_{44} - \frac{1}{2} e^{l\gamma+h} R = -\frac{\Lambda}{4M^3} e^{l\gamma+h}, \quad (8)$$

and

$$R_{a4} = -\frac{3}{2} k \sigma' h_{,a} = 0, \quad (9)$$

where the prime denotes derivative with respect to the additional y -coordinate. Eq.(9) implies two possibilities; either $\sigma' = 0$ ($\sigma = \text{const.}$) or $h_{,a} = 0$ ($h = \text{const.}$). In this letter we shall be interested in the 2nd possibility and we shall set $h = \text{const.} = 0$ so the line-element (4) can be written as,

$$ds^2 = e^{-2k\sigma(y)} g_{mn}(x) dx^m dx^n + e^{l\gamma(y)} dy^2. \quad (10)$$

Therefore the field equations (7) and (8) yield

$${}^4R_{ab} - \frac{1}{2}g_{ab} {}^4R = 3k e^{-2k\sigma - l\gamma}(\sigma'' - \frac{l}{2}\gamma'\sigma' - 2k\sigma'^2)g_{ab} - \frac{\Lambda}{4M^3}e^{-2k\sigma}g_{ab}, \quad (11)$$

and

$$\sigma'^2 = \frac{e^{l\gamma}}{12k^2}(e^{2k\sigma} {}^4R - \frac{\Lambda}{2M^3}), \quad (12)$$

respectively. ${}^4R_{ab}$ refers to the familiar four-dimensional Ricci tensor made out of the 4-d Christoffel symbols $\{^a_{bc}\} = \frac{1}{2}g^{an}(g_{bn,c} + g_{cn,b} - g_{bc,n})$ and ${}^4R = g^{mn} {}^4R_{mn}$ is the four-dimensional Ricci scalar. The 5-d and 4-d Ricci scalars are related through $R = e^{2k\sigma} {}^4R + 2k e^{-l\gamma}(4\sigma'' - 2l\gamma'\sigma' - 10k\sigma'^2)$. Combining the trace of Eq.(11) with Eq.(12) one obtains:

$$\sigma'' - \frac{l}{2}\gamma'\sigma' - k\sigma'^2 = \frac{\Lambda}{24kM^3}e^{l\gamma}. \quad (13)$$

For further simplification of our analysis let us set $g_{ab} = \eta_{ab}$ the usual four-dimensional Minkowski metric. In this case ${}^4R_{ab} = {}^4R = 0$ so, from Eq.(11), one gets

$$\sigma'' - \frac{l}{2}\gamma'\sigma' - 2k\sigma'^2 = \frac{\Lambda}{12kM^3}e^{l\gamma}. \quad (14)$$

Combining Eqs.(13) and (14) yields

$$\sigma'' = \frac{l}{2}\gamma'\sigma', \quad (15)$$

which, after integrating, gives

$$\sigma' = C e^{\frac{l}{2}\gamma}, \quad (16)$$

where C is some integration constant. If we put Eq.(15) into Eq.(13) or (14), we obtain

$$\sigma'^2 = -\frac{\Lambda}{24k^2M^3}e^{l\gamma}, \quad (17)$$

so the integration constant $C = \sqrt{-\frac{\Lambda}{24k^2M^3}}$. Eq.(17) (or Eq.(16)) makes sense only for negative $\Lambda < 0$ (i. e., $C^2 > 0$ and the constant C is real). This leads that our model five-dimensional spacetime is AdS_5 .

If we introduce a new coordinate through $\frac{d\sigma}{\sigma'} = dy$, it is encouraging noting that the line-element (10) can then be written as

$$ds^2 = e^{-2k|\sigma|}\eta_{mn}dx^m dx^n + r_c^2 d\sigma^2, \quad (18)$$

where $r_c = \sqrt{\frac{24k^2M^3}{-\Lambda}}$ and the orbifold symmetry $\sigma \rightarrow -\sigma$ has been taken into account. While deriving Eq.(18) we have used the following chain of equalities: $e^{l\gamma} dy^2 = \frac{e^{l\gamma}}{\sigma'^2} d\sigma^2 = \frac{24k^2M^3}{-\Lambda} d\sigma^2$. The line-element (18) exactly coincides with that of Randall and Sundrum if we set $\Lambda = -24k^2M^3$.

The most interesting feature of our set-up is contained in Eq.(17) (or Eq.(16)). Since both σ and γ are functions of the coordinate y , Eq.(17) can be written in the general form

$$\sigma'^2 + U(\sigma) = \mathcal{E}, \quad (19)$$

where $\mathcal{E}$ is an arbitrary positive constant and $U(\sigma)$ is an arbitrary function of σ . Then Eq.(19) implies that (see Eq.(17))

$$e^{l\gamma} = r_c^2 [\mathcal{E} - U(\sigma)]. \quad (20)$$

Therefore, any solution $\sigma(y)$ of the differential equation (19) in the form

$$\pm \int \frac{d\sigma}{\sqrt{\mathcal{E} - U(\sigma)}} = y + C_1, \quad (21)$$

implies, in virtue of Eq.(20), a solution $\gamma(y)$. This feature enables us some freedom in the choice of the function $U(\sigma)$. This is, precisely, the feature we shall exploit in searching for a mechanism of orbifold geometry stabilization. In fact, Eq.(19) may be given the following particle interpretation: it represents a scalar σ -particle with kinetic energy σ'^2 and total

energy $\mathcal{E}$, that moves along the y -direction in a potential $U(\sigma)$. In other words this can be put as follows. The derivative of Eq.(20) yields

$$l\gamma' = -\frac{dU}{d\sigma}\sigma' / (\mathcal{E} - U), \quad (22)$$

so, if we put Eq.(22) into Eq.(15) and, taking into account Eq.(19), one gets

$$\sigma'' = -\frac{1}{2} \frac{dU}{d\sigma}. \quad (23)$$

This equation of motion can also be obtained with the help of the variational principle from the action $S_y = \int dy \mathcal{L}[\sigma', \sigma]$, where $\mathcal{L}[\sigma', \sigma] = \sigma'^2 - U(\sigma)$.¹ Solutions to the equation of motion (23) that correspond to states of least energy are those with $\sigma = \sigma_i$ such that $U(\sigma_i)$ is a minimum. For these solutions the classical hamiltonian $H \sim \int dy [\sigma'^2 + U(\sigma)]$ is a minimum too. Therefore, solutions to Eq.(23) for which $\sigma = \sigma_i$ represent the ground σ -states of the system. These ground states stabilize the orbifold geometry in the sense that the points $\sigma = \sigma_i$ correspond to stable (ground) configurations of the scalar field σ , yielding that branes located at these points are stable against small perturbations of σ .

One instructive example is provided by the potential $U(\sigma) = \lambda\sigma^2$, where λ is an arbitrary constant. For positive $\lambda > 0$ Eq.(21) can be readily integrated to give

$$\sigma(y) = \sqrt{\frac{\mathcal{E}}{\lambda}} \sin \sqrt{\lambda} y, \quad (24)$$

where $-\frac{\pi}{2\sqrt{\lambda}} \leq y \leq \frac{\pi}{2\sqrt{\lambda}}$, so $-\sqrt{\frac{\mathcal{E}}{\lambda}} \leq \sigma \leq \sqrt{\frac{\mathcal{E}}{\lambda}}$ (we have set $C_1 = 0$). In this case the five-dimensional spacetime has two boundaries that are located at $\sigma = -\sqrt{\frac{\mathcal{E}}{\lambda}}$ and $\sigma = \sqrt{\frac{\mathcal{E}}{\lambda}}$ respectively. However these are not stable configurations in σ . The only stable configuration may be localized at $\sigma = 0$. For negative $\lambda < 0$, we would have, instead,

$$\sigma(y) = \sqrt{\frac{\mathcal{E}}{-\lambda}} \sinh \sqrt{-\lambda} y, \quad (25)$$

¹From Eq.(23) one sees that the potential should respect orbifold symmetry, i. e., $U(\sigma) = U(-\sigma)$

where now $-\infty \leq y \leq \infty$ and $-\infty \leq \sigma \leq \infty$ so, the σ -direction is unbounded. In this case there is no stable configuration supporting four-dimensional physics. We recall that this is just an instructive example.

A potential that supports the Randall-Sundrum mechanism for generating the mass hierarchy is the following

$$U(\sigma) = \lambda(\sigma_0^2 - \sigma^2)^2, \quad (26)$$

where $-\infty \leq \sigma \leq \infty$, i. e., we have an orbifold of AdS_5 geometry with infinite extent in the σ -direction. Following Ref. [5] we locate the Planck brane at the origin of the σ -coordinate $\sigma = 0$. The constant σ_0 is taken in such a way that $e^{-k|\sigma_0|} = TeV/M_{Pl}$ ($k \sim 10^{19} GeV$). For positive $\lambda > 0$, the potential $U(\sigma)$ in Eq.(26) has two minima at $\sigma = \sigma_0$ and $\sigma = -\sigma_0$ respectively. The origin $\sigma = 0$ is a local maximum. In other words, the stable ground σ -states are located at the minima $\sigma = \pm\sigma_0$.² In this case the electroweak TeV scale is generated from the Planck scale by spontaneous orbifold symmetry breaking. In fact, if the system was initially at $\sigma = 0$ (the Planck brane) it "rolls down" until the ground state at $\sigma = \sigma_0$ or at $\sigma = -\sigma_0$ is reached. Once the ground state is reached, say at $\sigma = -\sigma_0$, the orbifold symmetry inherent to the line-element (18) is not a symmetry of this ground state.

This mechanism also applies in the case of an arbitrary metric g_{ab} if we make the identification

$${}^4\Lambda \equiv \left(\frac{\Lambda}{4M^3} - 3ke^{-l\gamma}[\sigma'' - \frac{l}{2}\gamma'\sigma' - 2k\sigma'^2]\right)e^{-2k\sigma}, \quad (27)$$

where ${}^4\Lambda$ is a four-dimensional cosmological constant. In this case Eq.(11) can be written as

²For negative $\lambda < 0$, instead, this potential has two maxima at $\sigma = \pm\sigma_0$ and a local minimum at $\sigma = 0$. This theory has no bound σ -states since there may be tunneling from the region $|\sigma| \leq \sigma_0$ into the unbounded region $|\sigma| > \sigma_0$

$${}^4R_{ab} - \frac{1}{2} {}^4R = - {}^4\Lambda g_{ab}, \quad (28)$$

therefore, the requirement that the 4-d cosmological constant should be zero (${}^4\Lambda = 0$) enables our mechanism to work since, in this case, Eq.(14) and the subsequent set of equations (including Eq.(17)) hold true.

What we have proposed here is a very simple mechanism, literally a mechanism of spontaneous orbifold symmetry breaking, that allows stabilization of the orbifold geometry while solving the hierarchy problem a la Randall-Sundrum. However it is not addressed to give a realistic particle picture. A more realistic approach must contain matter sectors. This will be the subject of forthcoming work.

I acknowledge useful conversations with colleagues Rolando Cardenas and Rolando Bonal and MES of Cuba by financial support of this research.

REFERENCES

- [1] P. Horava and E. Witten, Nucl. Phys. B **460**, 506(1996); E. Witten, Nucl. Phys. B **471**, 135(1996); P. Horava and E. Witten, Nucl. Phys. B **475**, 94(1996).
- [2] J. E. Lidsey, D. Wands and E. J. Copeland, Phys. Rept. **337**, 343(2000).
- [3] A. Lukas, B. A. Ovrut, K. S. Stelle and D. Waldram, Phys. Rev. D **59**, 086001(1999); H. S. Reall, Phys. Rev. D **59**, 103506(1999).
- [4] L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 3370(1999); *ibid* **83**, 4690(1999).
- [5] J. Lykken and L. Randall, JHEP**0006**, 014(2000).